# **Elabscience**<sup>®</sup>

## Anti-2019-nCoV S-IgA Neutralizing Antibody

Catalog No.:	E-AB-V1027	Applications:	Neutralization, ELISA
Storage:	Store at -20°C , Avoid freeze / thaw cycles	Clone No.	8A5

Important Note: Centrifuge before opening to ensure complete recovery of vial contents.

### **Antigen Information**

## **Alternate Names** coronavirus s1, coronavirus s2, coronavirus spike, cov spike, ncov RBD, ncov s1, ncov s2, ncov spike, novel coronavirus RBD, novel coronavirus s1, novel coronavirus s2, novel coronavirus spike,RBD,S1,s2,Spike RBD Background Protein S (PROS1) is glycoprotein and expressed in many cell types supporting its reported involvement in multiple biological processes that include coagulation, apoptosis, cancer development and progression, and the innate immune response. Known receptors bind S1 are ACE2, angiotensinconverting enzyme 2, DPP4, CEACAM etc.. The spike (S) glycoprotein of coronaviruses is known to be essential in the binding of the virus to the host cell at the advent of the infection process. Most notable is severe acute respiratory syndrome (SARS). The severe acute respiratory syndrome-coronavirus (SARS-CoV) spike (S) glycoprotein alone can mediate the membrane fusion required for virus entry and cell fusion. It is also a major immunogen and a target for entry inhibitors. It's been reported that 2019-nCoV can infect the human respiratory epithelial cells through interaction with the human ACE2 receptor. The spike protein is a large type I transmembrane protein containing two subunits, S1 and S2. S1 mainly contains a receptor binding domain (RBD), which is responsible for recognizing the cell surface receptor. S2 contains basic elements needed for the membrane fusion. The S protein plays key parts in the induction of neutralizing-antibody and T-cell responses, as well as protective immunity.

## **Product Details**

Clone No.	8A5	
Isotype	Human IgA	
Immunogen	Recombinant 2019-nCoV S-trimer Protein (His Tag)[PKSR030489]	
Application	Neutralization, ELISA	
Reactivity	SARS-COV2	
Dilution	ELISA: 1:1,000-1:2,000	
Storage Buffer	PBS, pH 7.4, 10% glycerol	
Stability & Storage	Ships on ice packs. Store at -20°C	
Description	Antibody recognizes 2019-nCoV S Protein/ 2019-nCoV S-RBD Protein and block ACE-2 receptor	
	binding. The IgA fraction of the cell culture supernatant was purified by Protein A affinity	
	chromatography.	


#### For Research Use Only

Thank you for your recent purchase.

If you would like to learn more about antibodies, please visit www.elabscience.com Service for life science Applications:WB-Western Blot IHC-Immunohistochemistry IF-Immunofluorescence IP-Immunoprecipitation FC-Flow cytometry ChIP:Chromatin Immunoprecipitation

Focus on your research Service for life science

# **Elabscience**<sup>®</sup>



Immobilized Recombinant 2019nCoV S-trimer Protein (C-6His) (Cat#PKSR030489) at 5.0 ug/mL (100 uL/well) can bind Anti-2019- nCoV S-IgA Neutralizing Antibody (8A5), the EC50 is 397 ng/mL.

Immobilized Recombinant Recombinant 2019-nCoV S Protein RBD-SD1 (C-6His) (Cat#PKSR030477) at 5.0 ug/mL(100 uL/well) can bind Anti-2019- nCoV S-IgA Neutralizing Antibody (8A5), the EC50 is 416.4 ng/mL.

Anti-2019-nCoV S-IgA Neutralizing Antibody(8A5) can block Human ACE-2 Protein (Avi-His Tag)(Cat#PKSR030493) and 2019- nCoV S-trimer Protein (Cat#PKSR030489) interaction, the IC50 for this effect is 19.8 ng/mL.

### For Research Use Only

Thank you for your recent purchase.

Focus on your research Service for life science

If you would like to learn more about antibodies, please visit www.elabscience.com Service for life scient Applications:WB-Western Blot IHC-Immunohistochemistry IF-Immunofluorescence IP-Immunoprecipitation FC-Flow cytometry ChIP:Chromatin Immunoprecipitation