

## TBST Buffer, pH7.4 (10×)

Cat. No: E-BC-R335

Size: 100 mL/ 200 mL/ 500 mL

Cat	Products	100 mL	200 mL	500 mL	Storage
E-BC-R335	TBST Buffer, pH7.4 (10×)	100 mL	200 mL	500 mL	2~8°C

### Introduction

TBST buffer is used in Western Blot to wash away non-specific binding on the membrane, this buffer contains 0.1% Tween20. Tween20 is a non-ionic detergent that increases the buffer's ability to elute and reduce non-specific binding of antibodies. This product is supplied with 10× concentrate and easy for transportation.

### Instructions

This product is a 10× concentrate solution, dilute with Deionized water to 1× working solution before use. For example: take 100 mL of TBST buffer (10×), then add 900 mL of Deionized water and mix fully..

### 1 × TBST Buffer Components

145.4 mM NaCl, 10 mM Tris-base, 0.1% (v/v) Tween20, pH 7.4.

### Storage

Store at 2~8°C for 12 months.

### Cautions

1. If the product crystallizes, please dissolve in water bath at 37°C, and then dilute it after it is completely dissolved.
2. Please pay attention to sealed storage to avoid contamination
3. For your safety and health, please wear the lab coat and disposable gloves before the experiments.

### For Research Use Only