

2-step Plus Poly-HRP Anti Rabbit IgG Detection System (with DAB solution)

Cat. No: E-IR-R215

Size: 3 mL/ 6 mL/ 18 mL



Product Content

Cat	product	3 mL	6 mL	18 mL	Storage
E-IR-R215A	Normal Goat Blocking Buffer (Ready-to-Use)	3 mL	6 mL	18 mL	2~8 °C
E-IR-R215B	Polyperoxidase-anti-Rabbit IgG (Ready-to-Use)	3 mL	6 mL	18 mL	2~8 °C
E-IR-R215C	3% H ₂ O ₂	3 mL	6 mL	18 mL	2~8 °C
E-IR-R215D	DAB Concentrate (20×)	150 µL	300 µL	900 µL	2~8 °C
E-IR-R215E	DAB Substrate	3 mL	6 mL	18 mL	2~8 °C

Manual

One Copy

Introduction

2-step plus is a two-step immunohistochemical broad spectrum detection reagent. It polymerizes monovalent Fab fragments of secondary antibody and enzyme, which replaces the secondary antibody and tertiary antibody in traditional method, can directly amplify the binding signal of antibody-antigen. This method not only retains the specific binding ability of antibody with antigen, but also can effectively avoid space steric hindrance caused by excessive polymer molecules. Compared with the traditional SP three-step method, this kit has the characteristics of simple, rapid, high-sensitivity. This system abandons the using of biotin, so it can avoid background staining by endogenous biotin. It can be used in IHC, in which the primary antibody is monoclonal/polyclonal antibody derived from rabbit.

Diluent for DAB concentrated solution has been provided in this kit to avoid the influence of the different water acidity and alkalinity on the DAB Chromogenic Agent.

Sample dyeing

1. Dewax and hydrate the paraffin section.
2. Make thermal repair or digestion treatment to antigen of the tissue section if necessary according to antigen/antibody situation.
3. Incubate with E-IR-R215C (3% H₂O₂) for 10 min to eliminate endogenous peroxidase activity. Wash with PBS or TBS, 2 min×3 times.
4. Add E-IR-R215A (Normal Goat Blocking Buffer (Ready-to-Use)), incubate at 37 °C for 30 min. Shake off any excess liquid.
5. Add primary antibody (Rabbit-IgG) with proper dilution ratio, incubate at 20~37 °C for 1~2h or at 4 °C overnight (then rewarm at 37 °C for 30 min). Wash with PBS or TBS, 2 min×3 times. Dry the section with absorbent paper.
6. Add E-IR-R215B (Polyperoxidase-anti-Rabbit IgG), incubate at room temperature or 37 °C for 20 min. Wash with PBS or TBS, 2 min×3 times.
7. Add 1 drop (approximately 50 µL) of E-IR-R215D (DAB Concentrate) into each 1 mL of E-IR-R215E (DAB Substrate), mix fully and the mixed reagent is the DAB Working Solution. Prepare fresh solution before use and the prepared solution should be stored in the dark. Fresh prepared DAB Working Solution is valid within 4 hours and the unused solution must be abandoned
8. Take control of the DAB coloration period, the color of tan or brownish yellow is the positive signal. Avoid of excessive reaction.
9. Wash the section with deionized water terminate the chromogenic reaction, then operate the procedures of counterstaining, dehydrating, transparentizing and sealing

Storage

Store at 2~8 °C, shading light. Avoid of freezing. Valid for 12 months. The reagents are valid within 6 months after opening.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V1.3