


Tel:240-252-7368(USA) Fax: 240-252-7376(USA) techsupport@elabscience.com Website: www.elabscience.com

FGF2 Polyclonal Antibody


E-AB-40259 Catalog No. Reactivity M,R Storage Store at -20°C. Avoid freeze / thaw cycles. Host Rabbit **Applications IHC Isotype IgG**

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Images



Immunohistochemistry of paraffinembedded Rat brain using Fgf2 Polyclonal Antibody at dilution of 1:100



Immunohistochemistry of paraffinembedded Mouse brain using Fgf2 Polyclonal Antibody at dilution of 1:100

Immunogen Information

Recombinant Rat Fibroblast growth factor 2 protein **Immunogen**

GeneID 54250 P13109 **Swissprot**

Synonyms Basic fibroblast growth factor, Basic fibroblast growth

factor bFGF,HBGF 2

Product Information

Buffer PBS with 0.05% Proclin300 and 50% glycerol, pH7.4.

Purify Antigen Affinity Purification

Dilution IHC 1:100-1:200

Background

The protein encoded by this gene is a member of the fibroblast growth factor (FGF) family. FGF family members bind heparin and possess broad mitogenic and angiogenic activities. This protein has been implicated in diverse biological processes, such as limb and nervous system development, wound healing, and tumor growth. The mRNA for this gene contains multiple polyadenylation sites, and is alternatively translated from non-AUG (CUG) and AUG initiation codons, resulting in five different isoforms with distinct properties. The CUG-initiated isoforms are localized in the nucleus and are responsible for the intracrine effect, whereas, the AUG-initiated form is mostly cytosolic and is responsible for the paracrine and autocrine effects of this FGF.