

Goat Anti-Rat IgG(H+L)(Biotin conjugated)

Catalog No. E-AB-1032

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Product Information

Immunogen: Rat IgG

Purification Method: Affinity purification

Conjugation: Biotin

Concentration: 0.2 mg/mL(specific antibody purification≥99%),Biotin:Ab=20:1

Applications: IHC,WB,ELISA

Dilution: Immunohistochemistry: 1:100-200

Western Blotting: 1:300-10000(Enhanced chemiluminescent detection) Western Blotting: 1:200-400(DAB detection)

Direct ELISA: 1:5000-20000(TMB detection)

Optimal working dilutions must be determined by end user. Dilute before when it's needed, and the residual should be discarded.

Buffer: PBS with 50% glycerol, PH7.4

Storage: Store at -20°C. Avoid freeze / thaw cycles.

For Research Use Only

Thank you for your recent purchase.

If you would like to learn more about antibodies, please visit www.elabscience.com.

Focus on your research
Service for life science

Applications: Activ- Activation; Block- Blocking; Separation- Cell Separation ; Cell Sep-Neg- Cell Separation by Negative Selection; FA- Functional Assay; Neut- Neutralization; Stim- Stimulation; FCM- Flow Cytometry; ICFCM: Intracellular Staining for Flow Cytometry; WB- Western Blotting; IHC- Immunohistochemistry; IF- Immunofluorescence; IP- Immunoprecipitation