

Goat Anti-Rabbit IgG(H+L)(Rhodamine/TRITC conjugated)

Catalog No. E-AB-1053

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Product Information

Immunogen: Rabbit IgG

Purification Method: Affinity purification

Conjugation: TRITC

Concentration: 0.2 mg/mL,TRITC:Ab=4-6:1

Applications: IF

Dilution: Immunofluorescence: 1:20-100

Optimal working dilutions must be determined by end user. Dilute before when it's needed, and the residual should be discarded.

Amax: 550nm

E_{max}: 620nm

Buffer: PBS with 50% glycerol, PH7.4

Storage: Store at -20°C. Avoid freeze / thaw cycles. Protected from prolonged exposure to light.

For Research Use Only

Thank you for your recent purchase.

If you would like to learn more about antibodies, please visit www.elabscience.com.

Focus on your research
Service for life science

Applications: Activ- Activation; Block- Blocking; Separation- Cell Separation ; Cell Sep-Neg- Cell Separation by Negative Selection; FA- Functional Assay; Neut- Neutralization; Stim- Stimulation; FCM- Flow Cytometry; ICFM: Intracellular Staining for Flow Cytometry; WB- Western Blotting; IHC- Immunohistochemistry; IF- Immunofluorescence; IP- Immunoprecipitation