

## Rabbit Anti-Goat IgG(H+L)(Rhodamine/TRITC conjugated)

Catalog No. E-AB-1066

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.

### Product Information

**Immunogen:** Goat IgG

**Purification Method:** Affinity purification

**Conjugation:** TRITC

**Concentration:** 0.2 mg/mL,TRITC:Ab=4-6:1

**Applications:** IF

**Dilution:** Immunofluorescence: 1:20-100

Optimal working dilutions must be determined by end user. Dilute before when it's needed, and the residual should be discarded.

**Amax:** 550nm

**E<sub>max</sub>:** 620nm

**Buffer:** PBS with 50% glycerol, PH7.4

**Storage:** Store at -20°C. Avoid freeze / thaw cycles. Protected from prolonged exposure to light.

### For Research Use Only

Thank you for your recent purchase.

If you would like to learn more about antibodies, please visit [www.elabscience.com](http://www.elabscience.com).

Focus on your research  
Service for life science

Applications: Activ- Activation; Block- Blocking; Separation- Cell Separation ; Cell Sep-Neg- Cell Separation by Negative Selection; FA- Functional Assay; Neut- Neutralization; Stim- Stimulation; FCM- Flow Cytometry; ICFM: Intracellular Staining for Flow Cytometry; WB- Western Blotting; IHC- Immunohistochemistry; IF- Immunofluorescence; IP- Immunoprecipitation