


Tel:240-252-7368(USA) Fax: 240-252-7376(USA) techsupport@elabscience.com Website: www.elabscience.com

CDT1 Polyclonal Antibody


Catalog No.E-AB-17785ReactivityHStorageStore at -20°C. Avoid freeze / thaw cycles.HostRabbitApplicationsWB,IHC,ELISAIsotypeIgG

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Images



Western blot analysis of Human fetal brain tissue lysate using CDT1 Polyclonal Antibody at dilution of 1:400



Immunohistochemistry of paraffinembedded Human lung cancer tissue using CDT1 Polyclonal Antibody at dilution of 1:45(×200)



Immunohistochemistry of paraffinembedded Human liver cancer tissue using CDT1 Polyclonal Antibody at dilution of 1:45(×200)

Immunogen Information

Immunogen Synthetic peptide of human CDT1

Gene Accession NP112190 **Swissprot** Q9H211

Synonyms CDT 1,cdt1,CDT1,Retroviral integration

site1,Retroviral integration site2,RIS 2,RIS2

Product Information

Calculated MW 60 kDa

Observed MW Refer to figures

Buffer PBS with 0.05% NaN3 and 40% Glycerol,pH7.4

Purify Antigen affinity purification

Dilution WB 1:500-1:2000, IHC 1:30-1:150, ELISA

1:5000-1:10000

Background

The protein encoded by this gene is involved in the formation of the prereplication complex that is necessary for DNA replication. The encoded protein can bind geminin, which prevents replication and may function to prevent this protein from initiating replication at inappropriate origins. Phosphorylation of this protein by cyclin A-dependent kinases results in degradation of the protein.

For Research Use Only

Thank you for your recent purchase

If you would like to learn more about antibodies, please visit www.elabscience.com.

Focus on your research Service for life science