


Tel:240-252-7368(USA) Fax: 240-252-7376(USA) techsupport@elabscience.com Website: www.elabscience.com

HLA-C Polyclonal Antibody

Catalog No.E-AB-17922ReactivityHStorageStore at -20°C. Avoid freeze / thaw cycles.HostRabbitApplicationsIHC,ELISAIsotypeIgG

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Images



Immunohistochemistry of paraffinembedded Human colorectal cancer tissue using HLA-C Polyclonal Antibody at dilution of 1:25(×200)



Immunohistochemistry of paraffinembedded Human cervical cancer tissue using HLA-C Polyclonal Antibody at dilution of 1:25(×200)

Immunogen Information

Immunogen Synthetic peptide of human HLA-C

Gene Accession NP002108 **Swissprot** P10321

Synonyms HLA-JY3,D6S204,HLAC,Psoriasis Susceptibility

1,HLA-C Alpha Chain,PSORS1,MHC

Product Information

Buffer PBS with 0.05% NaN3 and 40% Glycerol,pH7.4

Purify Antigen affinity purification

Dilution IHC 1:25-1:100, ELISA 1:5000-1:10000

Background

HLA-C belongs to the HLA class I heavy chain paralogues. This class I molecule is a heterodimer consisting of a heavy chain and a light chain (beta-2 microglobulin). The heavy chain is anchored in the membrane. Class I molecules play a central role in the immune system by presenting peptides derived from endoplasmic reticulum lumen. They are expressed in nearly all cells. The heavy chain is approximately 45 kDa and its gene contains 8 exons. Exon one encodes the leader peptide, exons 2 and 3 encode the alpha1 and alpha2 domain, which both bind the peptide, exon 4 encodes the alpha3 domain, exon 5 encodes the transmembrane region, and exons 6 and 7 encode the cytoplasmic tail. Polymorphisms within exon 2 and exon 3 are responsible for the peptide binding specificity of each class one molecule. Typing for these polymorphisms is routinely done for bone marrow and kidney transplantation. Over one hundred HLA-C alleles have been described.