


Tel:240-252-7368(USA) Fax: 240-252-7376(USA) techsupport@elabscience.com Website: www.elabscience.com

KDM2A Polyclonal Antibody


Catalog No.E-AB-17949ReactivityH,MStorageStore at -20°C. Avoid freeze / thaw cycles.HostRabbitApplicationsIHC,ELISAIsotypeIgG

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Images



Immunohistochemistry of paraffinembedded Human lung cancer tissue using KDM2A Polyclonal Antibody at dilution of 1:55(×200)



Immunohistochemistry of paraffinembedded Human esophagus cancer tissue using KDM2A Polyclonal Antibody at dilution of 1:55(×200)

Immunogen Information

Immunogen Synthetic peptide of human KDM2A

Gene Accession NP036440 **Swissprot** Q9Y2K7

Synonyms FBL11,FBL7,FBXL11,JHDM1A,Lysine-specific

demethylase 2A

Product Information

Buffer PBS with 0.05% NaN3 and 40% Glycerol,pH7.4

Purify Antigen affinity purification

Dilution IHC 1:50-1:300, ELISA 1:5000-1:10000

Background

This gene encodes a member of the F-box protein family which is characterized by an approximately 40 amino acid motif, the F-box. The F-box proteins constitute one of the four subunits of ubiquitin protein ligase complex called SCFs (SKP1-cullin-F-box), which function in phosphorylation-dependent ubiquitination. The F-box proteins are divided into 3 classes: Fbws containing WD-40 domains, Fbls containing leucinerich repeats, and Fbxs containing either different protein-protein interaction modules or no recognizable motifs. The protein encoded by this gene belongs to the Fbls class and, in addition to an F-box, contains at least six highly degenerated leucine-rich repeats. This family member plays a role in epigenetic silencing. It nucleates at CpG islands and specifically demethylates both mono- and di-methylated lysine-36 of histone H3. Alternative splicing results in multiple transcript variants.