
# **Elabscience**®

## **TGM3 Polyclonal Antibody**


Catalog No.	E-AB-18250	Reactivity	Н
Storage	Store at -20°C. Avoid freeze / thaw cycles.	Host	Rabbit
Applications	WB,IHC,ELISA	Isotype	IgG

Note: Centrifuge before opening to ensure complete recovery of vial contents.

#### Images



Western blot analysis of Human tongue tissue lysate using TGM3 Polyclonal Antibody at dilution of 1:800



Immunohistochemistry of paraffinembedded Human liver cancer tissue using TGM3 Polyclonal Antibody at dilution of 1:55(×200)

#### **Immunogen Information**

Immunogen	Fusion protein of human TGM3
Gene Accession	BC109076
Swissprot	Q08188
Synonyms	TGase E,TGase-3,TGE,TGM 3,TGM3,TGM3,Transglutaminase E,TransglutaminaseE

### **Product Information**

Calculated MW	77 kDa
Observed MW	Refer to figures
Buffer	PBS with 0.05% NaN3 and 40% Glycerol,pH7.4
Purify	Antigen affinity purification
Dilution	WB 1:1000-1:5000, IHC 1:50-1:200, ELISA
	1:5000-1:10000

#### Background

Transglutaminases are enzymes that catalyze the crosslinking of proteins by epsilon-gamma glutamyl lysine isopeptide bonds. While the primary structure of transglutaminases is not conserved, they all have the same amino acid sequence at their active sites and their activity is calciumdependent. The protein encoded by this gene consists of two polypeptide chains activated from a single precursor protein by proteolysis. The encoded protein is involved the later stages of cell envelope formation in the epidermis and hair follicle.

For Research Use Only

Thank you for your recent purchase. If you would like to learn more about antibodies, please visit www.elabscience.com. Focus on your research Service for life science

Applications:WB-Western Blot IHC-Immunohistochemistry IF-Immunofluorescence IP-Immunoprecipitation FC-Flow cytometry ChIP-Chromatin Immunoprecipitation Reactivity: H-Human R-Rat M-Mouse Mk-Monkey Dg-Dog Ch-Chicken Hm-Hamster Rb-Rabbit Sh-Sheep Pg-Pig Z-Zebrafish X-Xenopus C-Cow.