


Tel:240-252-7368(USA) Fax: 240-252-7376(USA) techsupport@elabscience.com Website: www.elabscience.com

NDUFB11 Polyclonal Antibody

Catalog No.E-AB-18334ReactivityH,MStorageStore at -20°C. Avoid freeze / thaw cycles.HostRabbitApplicationsWB,ELISAIsotypeIgG

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Images



Western blot analysis of Hela HEPG2 A431 K562 and 231 cell Mouse liver tissue and Mouse brain tissue lysates using NDUFB11 Polyclonal Antibody at dilution of 1:1000

Immunogen Information

Immunogen Full length fusion protein

Gene Accession BC010665 **Swissprot** Q9NX14

Synonyms ESSS,NADH ubiquinone oxidoreductase ESSS

subunit, NDUFB 11, Neuronal protein

17.3,Np15,NP17.3,P17.3

Product Information

Calculated MW 17 kDa

Observed MW Refer to figures

Buffer PBS with 0.05% NaN3 and 40% Glycerol,pH7.4

Purify Antigen affinity purification

Dilution WB 1:1000-1:5000, ELISA 1:5000-1:10000

Background

Complex 1 (also known as NADH dehydrogenase) of the electron transport chain (respiratory chain) is an enzymatic complex that catalyzes the transfer of electrons from NADH to ubiquinone. Free energy from the reaction is conserved in the transfer of protons into the intermembrane space to create an electrochemical proton gradient, a driving force for ATP synthesis. Complex 1 is a complicated, multi-protein, L-shaped complex composed of at least 45 different subunits and located in the mitochondrial inner membrane. NDUFB11 (NADH dehydrogenase (ubiquinone) 1 beta subcomplex subunit 11), also known as ESSS, Np15, Np17.3 (neuronal protein 17.3) or p17.3, is a hydrophobic transmembrane protein belonging to the Complex I NDUFB11 subunit family. Ubiquitously expressed, NDUFB11 localizes to the inner membrane of the mitochondrion and functions as an accessory subunit of Complex I. The cAMP-dependent phosphorylation of NDUFB11 is important for the regulation of Complex I activity.

Focus on your research Service for life science