


Tel:240-252-7368(USA) Fax: 240-252-7376(USA) techsupport@elabscience.com Website: www.elabscience.com

# **MRPS18C Polyclonal Antibody**

Catalog No.E-AB-18355ReactivityH,MStorageStore at -20°C. Avoid freeze / thaw cycles.HostRabbitApplicationsWB,IHC,ELISAIsotypeIgG

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.

## **Images**



Western blot analysis of 231 Hela K562 cell and Human bladder carcinoma tissue using MRPS18C Polyclonal Antibody at dilution of 1:800



Immunohistochemistry of paraffinembedded Human tonsil tissue using MRPS18C Polyclonal Antibody at dilution of 1:25(×200)

## **Immunogen Information**

Immunogen Full length fusion protein

**Gene Accession** BC005186 **Swissprot** Q9Y3D5

Synonyms 28S ribosomal protein S18 1,MRP-

S18-1,Mrps18c,RT18C

#### **Product Information**

Calculated MW 16 kDa

**Observed MW** Refer to figures

**Buffer** PBS with 0.05% NaN3 and 40% Glycerol,pH7.4

**Purify** Antigen affinity purification

**Dilution** WB 1:1000-1:5000, IHC 1:25-1:100, ELISA

1:2000-1:10000

#### **Background**

Mammalian mitochondrial ribosomal proteins are encoded by nuclear genes and help in protein synthesis within the mitochondrion. Mitochondrial ribosomes (mitoribosomes) consist of a small 28S subunit and a large 39S subunit. They have an estimated 75% protein to rRNA composition compared to prokaryotic ribosomes, where this ratio is reversed. Another difference between mammalian mitoribosomes and prokaryotic ribosomes is that the latter contain a 5S rRNA. Among different species, the proteins comprising the mitoribosome differ greatly in sequence, and sometimes in biochemical properties, which prevents easy recognition by sequence homology.