


Tel:240-252-7368(USA) Fax: 240-252-7376(USA) techsupport@elabscience.com Website: www.elabscience.com

BHMT2 Polyclonal Antibody


Catalog No.E-AB-18539ReactivityH,M,RStorageStore at -20°C. Avoid freeze / thaw cycles.HostRabbitApplicationsWB,IHC,ELISAIsotypeIgG

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Images



Western blot analysis of Human fetal liver tissue and Human liver tissue lysates using BHMT2 Polyclonal Antibody at dilution of 1:800



Immunohistochemistry of paraffinembedded Human thyroid cancer tissue using BHMT2 Polyclonal Antibody at dilution of 1:30(×200)

Immunogen Information

Immunogen Fusion protein of human BHMT2

Gene Accession BC020665 **Swissprot** Q9H2M3

Synonyms Betaine homocysteine methyltransferase 2,SMM hcy

methyltransferase

Product Information

Calculated MW 40 kDa

Observed MW Refer to figures

Buffer PBS with 0.05% NaN3 and 40% Glycerol,pH7.4

Purify Antigen affinity purification

Dilution WB 1:1000-1:5000, IHC 1:30-1:150, ELISA

1:5000-1:10000

Background

Homocysteine is a sulfur-containing amino acid that plays a crucial role in methylation reactions. Transfer of the methyl group from betaine to homocysteine creates methionine, which donates the methyl group to methylate DNA, proteins, lipids, and other intracellular metabolites. The protein encoded by this gene is one of two methyl transferases that can catalyze the transfer of the methyl group from betaine to homocysteine. Anomalies in homocysteine metabolism have been implicated in disorders ranging from vascular disease to neural tube birth defects such as spina bifida. Alternatively spliced transcript variants encoding different isoforms have been found for this gene.