

Tel:240-252-7368(USA) Fax: 240-252-7376(USA) techsupport@elabscience.com Website: www.elabscience.com

# **C5AR1 Polyclonal Antibody**

Catalog No.E-AB-18573ReactivityHStorageStore at -20°C. Avoid freeze / thaw cycles.HostRabbitApplicationsIHC,ELISAIsotypeIgG

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.

## **Images**



Immunohistochemistry of paraffinembedded Human liver cancer tissue using C5AR1 Polyclonal Antibody at dilution of 1:50(×200)



Immunohistochemistry of paraffinembedded Human esophagus cancer tissue using C5AR1 Polyclonal Antibody at dilution of 1:50(×200)

## **Immunogen Information**

**Immunogen** Fusion protein of human C5AR1

**Gene Accession** BC008982 **Swissprot** P21730

**Synonyms** C5a anaphylatoxin

receptor,C5A,C5aR,C5AR,c5ar1,C5R1,CD88,CD88

antigen, Complement component 5 receptor 1

#### **Product Information**

**Buffer** PBS with 0.05% NaN3 and 40% Glycerol,pH7.4

**Purify** Antigen affinity purification

**Dilution** IHC 1:40-1:200, ELISA 1:5000-1:10000

#### **Background**

Receptor for the chemotactic and inflammatory peptide anaphylatoxin C5a (PubMed:1847994, PubMed:8182049, PubMed:7622471, PubMed:9553099, PubMed:10636859, PubMed:15153520). The ligand interacts with at least two sites on the receptor: a high-affinity site on the extracellular N-terminus, and a second site in the transmembrane region which activates downstream signaling events (PubMed:8182049, PubMed:7622471, PubMed:9553099). Receptor activation stimulates chemotaxis, granule enzyme release, intracellular calcium release and superoxide anion production (PubMed:10636859, PubMed:15153520).