


Tel:240-252-7368(USA) Fax: 240-252-7376(USA) techsupport@elabscience.com Website: www.elabscience.com

CD86 Polyclonal Antibody

E-AB-18593 Catalog No. Reactivity Η Storage Store at -20°C. Avoid freeze / thaw cycles. Rabbit Host **Applications** WB,IHC,ELISA **Isotype IgG**

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Images



Western blot analysis of Human lymph gland tissue and Jurkat cell lysates using CD86 Polyclonal Antibody at dilution of 1:450



Immunohistochemistry of paraffinembedded Human gastric cancer tissue using CD86 Polyclonal Antibody at dilution of $1:30(\times 200)$

Immunogen Information

Fusion protein of human CD86 **Immunogen**

BC040261 **Gene Accession** P42081 **Swissprot**

Synonyms Activation B7-2 antigen 3,T-lymphocyte activation

antigen CD86,TS/A-2

Product Information

Calculated MW 38 kDa

Observed MW Refer to figures

Buffer PBS with 0.05% NaN3 and 40% Glycerol,pH7.4

Purify Antigen affinity purification

Dilution WB 1:500-1:2000, IHC 1:25-1:100, ELISA

1:5000-1:10000

Background

This gene encodes a type I membrane protein that is a member of the immunoglobulin superfamily. This protein is expressed by antigenpresenting cells, and it is the ligand for two proteins at the cell surface of T cells, CD28 antigen and cytotoxic T-lymphocyte-associated protein 4. Binding of this protein with CD28 antigen is a costimulatory signal for activation of the T-cell. Binding of this protein with cytotoxic Tlymphocyte-associated protein 4 negatively regulates T-cell activation and diminishes the immune response. Alternative splicing results in several transcript variants encoding different isoforms.