


Tel:240-252-7368(USA) Fax: 240-252-7376(USA) techsupport@elabscience.com Website: www.elabscience.com

TBCB Polyclonal Antibody


E-AB-18621 Reactivity Catalog No. H,M Storage Store at -20°C. Avoid freeze / thaw cycles. Rabbit Host **Applications** WB,IHC,ELISA **Isotype IgG**

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Images



Western blot analysis of Human cerebrum tissue lysate using TBCB Polyclonal Antibody at dilution of 1:800



Immunohistochemistry of paraffinembedded Human gastric cancer tissue using TBCB Polyclonal Antibody at dilution of $1:25(\times 200)$

Immunogen Information

Full length fusion protein **Immunogen**

BC005969 **Gene Accession** O99426 **Swissprot**

Synonyms CG22,CKAPI,Cytoskeleton associated protein

1, Tubulin-specific chaperone B

Product Information

Calculated MW 27 kDa

Observed MW Refer to figures

Buffer PBS with 0.05% NaN3 and 40% Glycerol,pH7.4

Purify Antigen affinity purification

Dilution WB 1:500-1:2000, IHC 1:20-1:100, ELISA

1:5000-1:10000

Background

Microtubules, the primary component of the cytoskeletal network, are highly dynamic structures composed of α/β Tubulin heterodimers. Biosynthesis of functional microtubules involve the participation of several chaperones, termed Tubulin folding cofactors A (TBCA), B (TBCB), D (TBCD), E (TBCE) and C (TBCC), that act on folding intermediates downstream of the cytosolic chaperon, alternatively named TCP. TBCB (tubulin folding cofactor B), also known as CG22, CKAP1 or CKAPI, is a 244 amino acid cytoplasmic protein containing one CAP-Gly domain and in widely expressed. TBCB is involved in the regulation of tubulin heterodimer dissociation and may function as a negative regulator of axonal growth.