


Tel:240-252-7368(USA) Fax: 240-252-7376(USA) techsupport@elabscience.com Website: www.elabscience.com

UBXN1 Polyclonal Antibody


Catalog No.E-AB-18676ReactivityH,M,RStorageStore at -20°C. Avoid freeze / thaw cycles.HostRabbitApplicationsWB,IHC,ELISAIsotypeIgG

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Images



Western blot analysis of Mouse brain tissue Hela and HEPG2 cell lysates using UBXN1 Polyclonal Antibody at dilution of 1:500



Immunohistochemistry of paraffinembedded Human thyroid cancer tissue using UBXN1 Polyclonal Antibody at dilution of 1:35(×200)



Immunohistochemistry of paraffinembedded Human cervical cancer tissue using UBXN1 Polyclonal Antibody at dilution of 1:35(×200)

Immunogen Information

Immunogen Full length fusion protein

Gene Accession BC001372 **Swissprot** Q04323

Synonyms 2B28,MGC124706,MGC6696,mY33K,Protein

2B28,RGD1309471,SAKS1,UBXD10,UBXN1

Product Information

Calculated MW 33 kDa

Observed MW Refer to figures

Buffer PBS with 0.05% NaN3 and 40% Glycerol,pH7.4

Purify Antigen affinity purification

Dilution WB 1:500-1:2000, IHC 1:40-1:200, ELISA

1:5000-1:10000

Background

Ubiquitin-binding protein that interacts with the BRCA1-BARD1 heterodimer, and regulates its activity. Specifically binds 'Lys-6'-linked polyubiquitin chains. Interaction with autoubiquitinated BRCA1, leads to inhibit the E3 ubiquitin-protein ligase activity of the BRCA1-BARD1 heterodimer. Component of a complex required to couple deglycosylation and proteasome-mediated degradation of misfolded proteins in the endoplasmic reticulum that are retrotranslocated in the cytosol.

For Research Use Only

Thank you for your recent purchase

If you would like to learn more about antibodies, please visit www.elabscience.com.

Focus on your research Service for life science