


Tel:240-252-7368(USA) Fax: 240-252-7376(USA) techsupport@elabscience.com Website: www.elabscience.com

ATP5L Polyclonal Antibody

Catalog No. E-AB-18764 Reactivity H,M Store at -20°C. Avoid freeze / thaw cycles. Rabbit **Storage** Host **Applications** WB,ELISA **Isotype IgG**

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Images



Western blot analysis of Human fetal liver tissue Hela cell HEPG2 cell lysates using ATP5L Polyclonal Antibody at dilution of 1:500

Immunogen Information

Fusion protein of human ATP5L **Immunogen**

Gene Accession BC015128 O75964 **Swissprot**

Synonyms ATP synthase subunit g,ATP

synthase, ATP5L, ATP5L, ATPase subunit

g,mitochondrial

Product Information

Calculated MW 11 kDa

Observed MW Refer to figures

Buffer PBS with 0.05% NaN3 and 40% Glycerol,pH7.4

Purify Antigen affinity purification

Dilution WB 1:500-1:2000, ELISA 1:5000-1:10000

Background

Mitochondrial ATP synthase catalyzes ATP synthesis, utilizing an electrochemical gradient of protons across the inner membrane during oxidative phosphorylation. It is composed of two linked multi-subunit complexes: the soluble catalytic core, F1, and the membrane-spanning component, Fo, which comprises the proton channel. The F1 complex consists of 5 different subunits (alpha, beta, gamma, delta, and epsilon) assembled in a ratio of 3 alpha, 3 beta, and a single representative of the other 3. The Fo seems to have nine subunits (a, b, c, d, e, f, g, F6 and 8). This gene encodes the g subunit of the Fo complex. Alternative splicing results in multiple transcript variants.