


Tel:240-252-7368(USA) Fax: 240-252-7376(USA) techsupport@elabscience.com Website: www.elabscience.com

# **HBG1/HBG2 Polyclonal Antibody**


Catalog No. E-AB-18774 Reactivity Η Storage Store at -20°C. Avoid freeze / thaw cycles. Rabbit Host **Applications** WB,IHC,ELISA **Isotype IgG** 

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.

## **Images**



Western blot analysis of Human plasma solution using HBG1:HBG2 Polyclonal Antibody at dilution of 1:250



Immunohistochemistry of paraffinembedded Human thyroid cancer tissue using HBG1:HBG2 Polyclonal Antibody at dilution of  $1:70(\times 200)$ 

## **Immunogen Information**

Fusion protein of human HBG1/HBG2 **Immunogen** 

BC010914 **Gene Accession** P69891,P69892 **Swissprot** 

**Synonyms** A gamma globin, HBG 1, HBG

2,HBG1,HBG2,HBGA,HBGR,HSGGL

1,HSGGL1,PRO2979

#### **Product Information**

Calculated MW 16 kDa

**Observed MW** Refer to figures

**Buffer** PBS with 0.05% NaN3 and 40% Glycerol,pH7.4

**Purify** Antigen affinity purification

Dilution WB 1:500-1:2000, IHC 1:40-1:200, ELISA

1:5000-1:10000

#### **Background**

The gamma globin genes (HBG1 and HBG2) are normally expressed in the fetal liver, spleen and bone marrow. Two gamma chains together with two alpha chains constitute fetal hemoglobin (HbF) which is normally replaced by adult hemoglobin (HbA) at birth. In some beta-thalassemias and related conditions, gamma chain production continues into adulthood. The two types of gamma chains differ at residue 136 where glycine is found in the G-gamma product (HBG2) and alanine is found in the Agamma product (HBG1). The former is predominant at birth. The order of the genes in the beta-globin cluster is: 5'-epsilon -- gamma-G -- gamma-A -- delta -- beta--3'.