


CD48 Polyclonal Antibody


| | | | |
|---------------------|---|-------------------|--------|
| Catalog No. | E-AB-18799 | Reactivity | H |
| Storage | Store at -20°C. Avoid freeze / thaw cycles. | Host | Rabbit |
| Applications | WB,IHC,ELISA | Isotype | IgG |

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Images



Western blot analysis of 231 K562 Jurkat and Raji cell lysates using CD48 Polyclonal Antibody at dilution of 1:550



Immunohistochemistry of paraffin-embedded Human tonsil tissue using CD48 Polyclonal Antibody at dilution of 1:95(×200)

Immunogen Information

| | |
|-----------------------|--|
| Immunogen | Fusion protein of human CD48 |
| Gene Accession | BC016182 |
| Swissprot | P09326 |
| Synonyms | Antigen CD48,BCM1,BLAST 1,BLAST,BLAST1,CD 48,CD48,mCD48,MEM 102,MEM-102,SLAMF2,TCT.1 |

Product Information

| | |
|----------------------|--|
| Calculated MW | 28 kDa |
| Observed MW | Refer to figures |
| Buffer | PBS with 0.05% NaN ₃ and 40% Glycerol,pH7.4 |
| Purify | Antigen affinity purification |
| Dilution | WB 1:500-1:2000, IHC 1:50-1:300, ELISA 1:5000-1:10000 |

Background

This gene encodes a member of the CD2 subfamily of immunoglobulin-like receptors which includes SLAM (signaling lymphocyte activation molecules) proteins. The encoded protein is found on the surface of lymphocytes and other immune cells, dendritic cells and endothelial cells, and participates in activation and differentiation pathways in these cells. The encoded protein does not have a transmembrane domain, however, but is held at the cell surface by a GPI anchor via a C-terminal domain which maybe cleaved to yield a soluble form of the receptor. Multiple transcript variants encoding different isoforms have been found for this gene.

For Research Use Only

Thank you for your recent purchase.
 If you would like to learn more about antibodies, please visit www.elabscience.com.

**Focus on your research
 Service for life science**

Applications:WB-Western Blot IHC-Immunohistochemistry IF-Immunofluorescence IP-Immunoprecipitation FC-Flow cytometry ChIP-Chromatin Immunoprecipitation Reactivity: H-Human R-Rat M-Mouse Mk-Monkey Dg-Dog Ch-Chicken Hm-Hamster Rb-Rabbit Sh-Sheep Pg-Pig Z-Zebrafish X-Xenopus C-Cow.