


Tel:240-252-7368(USA) Fax: 240-252-7376(USA) techsupport@elabscience.com Website: www.elabscience.com

# **HLA-DRB4 Polyclonal Antibody**


Catalog No.E-AB-18812ReactivityHStorageStore at -20°C. Avoid freeze / thaw cycles.HostRabbitApplicationsWB,IHC,ELISAIsotypeIgG

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.

## **Images**



Western blot analysis of Raji cell lysate using HLA-DRB4 Polyclonal Antibody at dilution of 1:450



Immunohistochemistry of paraffinembedded Human tonsil tissue using HLA-DRB4 Polyclonal Antibody at dilution of 1:60(×200)

## **Immunogen Information**

Immunogen Fusion protein of human HLA-DRB4

**Gene Accession** BC005312 **Swissprot** P13762

**Synonyms** DR 4,DR beta 4 chain,DRB4,DRB4,HLA-

DRB4,MHC class2 antigen ,MHC HLA DR-beta

chain

#### **Product Information**

Calculated MW 30 kDa

**Observed MW** Refer to figures

**Buffer** PBS with 0.05% NaN3 and 40% Glycerol,pH7.4

**Purify** Antigen affinity purification

**Dilution** WB 1:500-1:2000, IHC 1:20-1:100, ELISA

1:5000-1:10000

#### **Background**

HLA-DRB4 belongs to the HLA class II beta chain paralogues. This class II molecule is a heterodimer consisting of an alpha (DRA) and a beta (DRB) chain, both anchored in the membrane. It plays a central role in the immune system by presenting peptides derived from extracellular proteins. Class II molecules are expressed in antigen presenting cells (APC: B lymphocytes, dendritic cells, macrophages). The beta chain is approximately 26-28 kDa and its gene contains 6 exons. Exon one encodes the leader peptide, exons 2 and 3 encode the two extracellular domains, exon 4 encodes the transmembrane domain and exon 5 encodes the cytoplasmic tail. Within the DR molecule the beta chain contains all the polymorphisms specifying the peptide binding specificities. Typing for these polymorphisms is routinely done for bone marrow and kidney transplantation. DRB1 is expressed at a level five times higher than its paralogues DRB3, DRB4 and DRB5. The presence of DRB4 is linked with allelic variants of DRB1, otherwise it is omitted. There are 4 related pseudogenes: DRB2, DRB6, DRB7, DRB8 and DRB9.

For Research Use Only

Thank you for your recent purchase

If you would like to learn more about antibodies, please visit www.elabscience.com.

Focus on your research Service for life science