


Tel:240-252-7368(USA) Fax: 240-252-7376(USA) techsupport@elabscience.com Website: www.elabscience.com

PNRC2 Polyclonal Antibody


Catalog No.E-AB-18895ReactivityH,M,RStorageStore at -20°C. Avoid freeze / thaw cycles.HostRabbitApplicationsIHC,ELISAIsotypeIgG

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Images



Immunohistochemistry of paraffinembedded Human esophagus cancer tissue using PNRC2 Polyclonal Antibody at dilution of 1:30(×200)



Immunohistochemistry of paraffinembedded Human liver cancer tissue using PNRC2 Polyclonal Antibody at dilution of 1:30(×200)

Immunogen Information

Immunogen Fusion protein of human PNRC2

Gene Accession BC001959 **Swissprot** Q9NPJ4

Synonyms FLJ20312,MGC11707,MGC93828,MGC99541,pnrc2

,PNRC2,RP11-4M23.5,RP23-161N17.3

Product Information

Buffer PBS with 0.05% NaN3 and 40% Glycerol,pH7.4

Purify Antigen affinity purification

Dilution IHC 1:30-1:150, ELISA 1:5000-1:10000

Background

Involved in nonsense-mediated mRNA decay (NMD) by acting as a bridge between the mRNA decapping complex and the NMD machinery (PubMed:19150429). May act by targeting the NMD machinery to the P-body and recruiting the decapping machinery to aberrant mRNAs (PubMed:19150429). Required for UPF1/RENT1 localization to the P-body (PubMed:19150429). Plays a role in glucocorticoid receptor-mediated mRNA degradation by interacting with the glucocorticoid receptor NR3C1 in a ligand-dependent manner when it is bound to the 5' UTR of target mRNAs and recruiting the RNA helicase UPF1 and the mRNA-decapping enzyme DCP1A, leading to RNA decay (PubMed:25775514). Also acts as a nuclear receptor coactivator (PubMed:11574675). May play a role in controlling the energy balance between energy storage and energy expenditure (By similarity).