


Tel:240-252-7368(USA) Fax: 240-252-7376(USA) techsupport@elabscience.com Website: www.elabscience.com

NKD2 Polyclonal Antibody

Catalog No.E-AB-18970ReactivityH,MStorageStore at -20°C. Avoid freeze / thaw cycles.HostRabbitApplicationsWB,IHC,ELISAIsotypeIgG

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Images



Western blot analysis of Mouse Pancreas tissue lysate using NKD2 Polyclonal Antibody at dilution of 1:350



Immunohistochemistry of paraffinembedded Human tonsil tissue using NKD2 Polyclonal Antibody at dilution of 1:50(×200)



Immunohistochemistry of paraffinembedded Human prost ate cancer tissue using NKD2 Polyclonal Antibody at dilution of 1:50(×200)

Immunogen Information

Immunogen Fusion protein of human NKD2

Gene Accession BC012176 **Swissprot** Q969F2

Synonyms hNkd2,Naked 2,Naked-2,nkd2,NKD2,Protein naked

cuticle homolog 2

Product Information

Calculated MW 50 kDa

Observed MW Refer to figures

Buffer PBS with 0.05% NaN3 and 40% Glycerol,pH7.4

Purify Antigen affinity purification

Dilution WB 1:500-1:2000, IHC 1:40-1:200, ELISA

1:5000-1:10000

Background

This gene encodes a member of a family of proteins that function as negative regulators of Wnt receptor signaling through interaction with Dishevelled family members. The encoded protein participates in the delivery of transforming growth factor alpha-containing vesicles to the cell membrane. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene.

For Research Use Only

Thank you for your recent purchase

If you would like to learn more about antibodies, please visit www.elabscience.com.

Focus on your research Service for life science