


Tel:240-252-7368(USA) Fax: 240-252-7376(USA) techsupport@elabscience.com Website: www.elabscience.com

# **ECH1 Polyclonal Antibody**

E-AB-18979 Catalog No. Reactivity H,R Storage Store at -20°C. Avoid freeze / thaw cycles. Rabbit Host **Applications** WB,IHC,ELISA **Isotype IgG** 

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.

## **Images**



Western blot analysis of 293T HepG2 and A549 cell lysates using ECH1 Polyclonal Antibody at dilution of 1:200



Immunohistochemistry of paraffinembedded Human tonsil tissue using ECH1 Polyclonal Antibody at dilution of  $1:50(\times 200)$ 

## **Immunogen Information**

Fusion protein of human ECH1 **Immunogen** 

BC011792 **Gene Accession** O13011 **Swissprot** 

**Synonyms** ECH 1.Ech1.ECH1.HPXEL.mitochondrial,MS1034.P

eroxisomal enoyl CoA hydratase 1

#### **Product Information**

Calculated MW 36 kDa

**Observed MW** Refer to figures

**Buffer** PBS with 0.05% NaN3 and 40% Glycerol,pH7.4

**Purify** Antigen affinity purification

Dilution WB 1:500-1:2000, IHC 1:50-1:300, ELISA

1:5000-1:10000

### **Background**

This gene encodes a member of the hydratase/isomerase superfamily. The gene product shows high sequence similarity to enoyl-coenzyme A (CoA) hydratases of several species, particularly within a conserved domain characteristic of these proteins. The encoded protein, which contains a Cterminal peroxisomal targeting sequence, localizes to the peroxisome. The rat ortholog, which localizes to the matrix of both the peroxisome and mitochondria, can isomerize 3-trans,5-cis-dienoyl-CoA to 2-trans,4-transdienovl-CoA, indicating that it is a delta3,5-delta2,4-dienovl-CoA isomerase. This enzyme functions in the auxiliary step of the fatty acid beta-oxidation pathway. Expression of the rat gene is induced by peroxisome proliferators.