


Tel:240-252-7368(USA) Fax: 240-252-7376(USA) techsupport@elabscience.com Website: www.elabscience.com

# **ELOC Polyclonal Antibody**


Catalog No.E-AB-19103ReactivityH,M,RStorageStore at -20°C. Avoid freeze / thaw cycles.HostRabbitApplicationsIHC,ELISAIsotypeIgG

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.

# **Images**



Immunohistochemistry of paraffinembedded Human breast cancer tissue using ELOC Polyclonal Antibody at dilution of 1:55(×200)



Immunohistochemistry of paraffinembedded Human ovarian cancer tissue using ELOC Polyclonal Antibody at dilution of 1:55(×200)

# **Immunogen Information**

Immunogen Fusion protein of human ELOC

**Gene Accession** BC013809 **Swissprot** Q15369

**Synonyms** Transcription Elongation Factor B Subunit 1, Elongin

15 KDa Subunit, SIII P15.

#### **Product Information**

**Buffer** PBS with 0.05% NaN3 and 40% Glycerol,pH7.4

**Purify** Antigen affinity purification

**Dilution** IHC 1:50-1:300, ELISA 1:5000-1:10000

# **Background**

This gene encodes the protein elongin C, which is a subunit of the transcription factor B (SIII) complex. The SIII complex is composed of elongins A/A2, B and C. It activates elongation by RNA polymerase II by suppressing transient pausing of the polymerase at many sites within transcription units. Elongin A functions as the transcriptionally active component of the SIII complex, whereas elongins B and C are regulatory subunits. Elongin A2 is specifically expressed in the testis, and capable of forming a stable complex with elongins B and C. The von Hippel-Lindau tumor suppressor protein binds to elongins B and C, and thereby inhibits transcription elongation. Multiple alternatively spliced transcript variants encoding two distinct isoforms have been identified.