


Tel:240-252-7368(USA) Fax: 240-252-7376(USA) techsupport@elabscience.com Website: www.elabscience.com

PSMA6 Polyclonal Antibody


Catalog No.E-AB-19189ReactivityH,M,RStorageStore at -20°C. Avoid freeze / thaw cycles.HostRabbitApplicationsWB,IHC,ELISAIsotypeIgG

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Images



Western blot analysis of 293T cell lysates using PSMA6 Polyclonal Antibody at dilution of 1:1000



Immunohistochemistry of paraffinembedded Human cervical cancer tissue using PSMA6 Polyclonal Antibody at dilution of 1:90(×200)

Immunogen Information

Immunogen Fusion protein of human PSMA6

Gene Accession BC002979 **Swissprot** P60900

Synonyms Proteasome subunit iota,PSA6,PSMA 6,PSMA6

Product Information

Calculated MW 27 kDa

Observed MW Refer to figures

Buffer PBS with 0.05% NaN3 and 40% Glycerol,pH7.4

Purify Antigen affinity purification

Dilution WB 1:1000-1:5000, IHC 1:100-1:300, ELISA

1:5000-1:10000

Background

The proteasome is a multicatalytic proteinase complex with a highly ordered ring-shaped 20S core structure. The core structure is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. This gene encodes a member of the peptidase T1A family, that is a 20S core alpha subunit. Multiple transcript variants encoding several different isoforms have been found for this gene. A pseudogene has been identified on the Y chromosome.