


Tel:240-252-7368(USA) Fax: 240-252-7376(USA) techsupport@elabscience.com Website: www.elabscience.com

ASIC3 Polyclonal Antibody

Catalog No.E-AB-19330ReactivityHStorageStore at -20°C. Avoid freeze / thaw cycles.HostRabbitApplicationsIHC,ELISAIsotypeIgG

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Images



Immunohistochemistry of paraffinembedded Human colorectal cancer tissue using ASIC3 Polyclonal Antibody at dilution of 1:40(×200)

Immunogen Information

Immunogen Synthetic peptide of human ASIC3

Gene Accession NP004760 **Swissprot** Q9UHC3

Synonyms ASIC 3,ASIC3,DRASIC,hASIC 3,hASIC3,hTNaC

1,hTNaC1,TNaC1,TNaC1

Product Information

Buffer PBS with 0.05% NaN3 and 40% Glycerol,pH7.4

Purify Antigen affinity purification

Dilution IHC 1:30-1:150, ELISA 1:2000-1:10000

Background

This gene encodes a member of the degenerin/epithelial sodium channel (DEG/ENaC) superfamily. The members of this family are amiloridesensitive sodium channels that contain intracellular N and C termini, two hydrophobic transmembrane regions, and a large extracellular loop, which has many cysteine residues with conserved spacing. The member encoded by this gene is an acid sensor and may play an important role in the detection of lasting pH changes. In addition, a heteromeric association between this member and acid-sensing (proton-gated) ion channel 2 has been observed as proton-gated channels sensitive to gadolinium. Alternatively spliced transcript variants have been described.