


Tel:240-252-7368(USA) Fax: 240-252-7376(USA) techsupport@elabscience.com Website: www.elabscience.com

CDKN2A Polyclonal Antibody


Catalog No.E-AB-19334ReactivityHStorageStore at -20°C. Avoid freeze / thaw cycles.HostRabbitApplicationsIHC,ELISAIsotypeIgG

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Images



Immunohistochemistry of paraffinembedded Human gasrtic cancer tissue using CDKN2A Polyclonal Antibody at dilution of 1:50(×200)



Immunohistochemistry of paraffinembedded Human thyroid cancer tissue using CDKN2A Polyclonal Antibody at dilution of 1:50(×200)

Immunogen Information

Immunogen Synthetic peptide of human CDKN2A

Gene Accession NP000068 **Swissprot** P42771

Synonyms CCM2,CDK4 inhibitor p16

INK4,CDK4I,CDKN2,CDKN2A,Cell cycle negative regulator beta,CMM2,p14,p16,P16INK4,p16INK4a,p

19,p19Arf,TP16

Product Information

Buffer PBS with 0.05% NaN3 and 40% Glycerol,pH7.4

Purify Antigen affinity purification

Dilution IHC 1:50-1:200, ELISA 1:1000-1:5000

Background

This gene generates several transcript variants which differ in their first exons. At least three alternatively spliced variants encoding distinct proteins have been reported, two of which encode structurally related isoforms known to function as inhibitors of CDK4 kinase. The remaining transcript includes an alternate first exon located 20 Kb upstream of the remainder of the gene; this transcript contains an alternate open reading frame (ARF) that specifies a protein which is structurally unrelated to the products of the other variants. This ARF product functions as a stabilizer of the tumor suppressor protein p53 as it can interact with, and sequester, the E3 ubiquitin-protein ligase MDM2, a protein responsible for the degradation of p53.