


Tel:240-252-7368(USA) Fax: 240-252-7376(USA) techsupport@elabscience.com Website: www.elabscience.com

HAS1 Polyclonal Antibody


Catalog No.E-AB-19341ReactivityH,MStorageStore at -20°C. Avoid freeze / thaw cycles.HostRabbitApplicationsIHC,ELISAIsotypeIgG

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Images



Immunohistochemistry of paraffinembedded Human ovarian cancer tissue using HAS1 Polyclonal Antibody at dilution of 1:70(×200)



Immunohistochemistry of paraffinembedded Human lung cancer tissue using HAS1 Polyclonal Antibody at dilution of 1:70(×200)

Immunogen Information

Immunogen Synthetic peptide of human HAS1

Gene Accession NP001514 **Swissprot** Q92839

Synonyms HA synthase 1,HAS,HAS1,HuHAS1,Hyaluronan

synthase 1, Hyaluronate synthase 1, Hyaluronic acid

synthase 1,HYAS1

Product Information

Buffer PBS with 0.05% NaN3 and 40% Glycerol,pH7.4

Purify Antigen affinity purification

Dilution IHC 1:50-1:200, ELISA 1:5000-1:10000

Background

Hyaluronan or hyaluronic acid (HA) is a high molecular weight unbranched polysaccharide synthesized by a wide variety of organisms from bacteria to mammals, and is a constituent of the extracellular matrix. It consists of alternating glucuronic acid and N-acetylglucosamine residues that are linked by beta-1-3 and beta-1-4 glycosidic bonds. HA is synthesized by membrane-bound synthase at the inner surface of the plasma membrane, and the chains are extruded through pore-like structures into the extracellular space. It serves a variety of functions, including space filling, lubrication of joints, and provision of a matrix through which cells can migrate. HA is actively produced during wound healing and tissue repair to provide a framework for ingrowth of blood vessels and fibroblasts. Changes in the serum concentration of HA are associated with inflammatory and degenerative arthropathies such as rheumatoid arthritis.