


Tel:240-252-7368(USA) Fax: 240-252-7376(USA) techsupport@elabscience.com Website: www.elabscience.com

# **RGR Polyclonal Antibody**

Catalog No.E-AB-19982ReactivityH,MStorageStore at -20°C. Avoid freeze / thaw cycles.HostRabbitApplicationsWB,IHC,ELISAIsotypeIgG

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.

# **Images**



Western blot analysis of Mouse liver tissue lysate using RGR Polyclonal Antibody at dilution of 1:300



Immunohistochemistry of paraffinembedded Human tonsil tissue using RGR Polyclonal Antibody at dilution of 1:25(×200)

# **Immunogen Information**

Immunogen Synthetic peptide of human RGR

Gene Accession NP001012738

Swissprot P47804

**Synonyms** hRGR,Ral GDS related protein,Retinal G protein

coupled receptor,RGR-opsin,RP44,RPE retinal G

protein coupled receptor

#### **Product Information**

Calculated MW 32 kDa

**Observed MW** Refer to figures

**Buffer** PBS with 0.05% NaN3 and 40% Glycerol,pH7.4

**Purify** Antigen affinity purification

**Dilution** WB 1:500-1:2000, IHC 1:30-1:150, ELISA

1:5000-1:10000

# **Background**

This gene encodes a putative retinal G-protein coupled receptor. The gene is a member of the opsin subfamily of the 7 transmembrane, G-protein coupled receptor 1 family. Like other opsins which bind retinaldehyde, it contains a conserved lysine residue in the seventh transmembrane domain. The protein acts as a photoisomerase to catalyze the conversion of all-trans-retinal to 11-cis-retinal. The reverse isomerization occurs with rhodopsin in retinal photoreceptor cells. The protein is exclusively expressed in tissue adjacent to retinal photoreceptor cells, the retinal pigment epithelium and Mueller cells. This gene may be associated with autosomal recessive and autosomal dominant retinitis pigmentosa (arRP and adRP, respectively). Alternative splicing results in multiple transcript variants encoding different isoforms.