


Tel:240-252-7368(USA) Fax: 240-252-7376(USA) techsupport@elabscience.com Website: www.elabscience.com

ITPR2 Polyclonal Antibody


Catalog No.E-AB-52051ReactivityH,M,RStorageStore at -20°C. Avoid freeze / thaw cycles.HostRabbitApplicationsIHC,ELISAIsotypeIgG

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Images



Immunohistochemistry of paraffinembedded Human tonsil tissue using ITPR2 Polyclonal Antibody at dilution of 1:60(×200)



Immunohistochemistry of paraffinembedded Human esophagus cancer tissue using ITPR2 Polyclonal Antibody at dilution of 1:60(×200)

Immunogen Information

Immunogen Synthetic peptide of human ITPR2

Gene Accession NP002214 **Swissprot** Q14571

Synonyms InsP3 R2,InsP3R2,IP3 R2,IP3 receptor,IP3R

2,IP3R2,ITPR 2,Itpr2,ITPR2

Product Information

Buffer PBS with 0.05% NaN3 and 40% Glycerol,pH7.4

Purify Antigen affinity purification

Dilution IHC 1:50-1:300, ELISA 1:5000-1:10000

Background

Inositol 1,4,5-triphosphate (IP3) functions as a second messenger for a myriad of extracellular stimuli including hormones, growth factors and neurotransmitters. Receptor tyrosine kinases indirectly increase the intracellular levels of IP3 through the activation of phospholipases such as phospholipase C (PLC), which convert phosphatidylinositol-4,5 bisphosphate into IP3 and diacylglycerol (DAG). The inositol 1,4,5-triphosphate receptor, IP3R, acts as an inositol triphosphate (IP3)-gated calcium release channel in a variety of cell types. Three IP3 receptor subtypes have been described and are designated IP3R-I, IP3R-II and IP3R-III. IP3R-I is the predominant IP3R subtype expressed in neuronal tissues and the central nervous system, but is also expressed at high levels in the liver.