

Tel:240-252-7368(USA) Fax: 240-252-7376(USA) techsupport@elabscience.com Website: www.elabscience.com

beta Tubulin Monoclonal Antibody


Catalog No. E-AB-20033 Reactivity H,M,R,Mk,Ch,Dg,Hm,Rb,

Sh,Insect,Yeast

StorageStore at -20°C. Avoid freeze / thaw cycles.HostMouseApplicationsWB,IHC-p,IFIsotypeIgG

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Images



Western Blot analysis of A549, Rat brain, Mouse brain, Chicken lung, Rabbit testis, Sheep muscle using beta Tubulin Monoclonal Antibody at dilution of 1:5000.



Immunohistochemistry of paraffinembedded Mouse testis tissue using beta Tubulin Monoclonal Antibody at dilution of 1:200.



Immunofluorescence analysis of Hela tissue using beta Tubulin Monoclonal Antibody at dilution of 1:100.

Immunogen Information

Immunogen Synthetic Peptide

Swissprot Q13509

Synonyms TUBB3,TUBB4,Tubulin beta-3 chain,Tubulin beta-4

chain, Tubulin beta-III

Product Information

Calculated MW 50kDa
Observed MW 55kDa

Buffer PBS with 0.02% sodium azide and 50% glycerol pH

7.4.

Purify Protein A purification

Clone No. Clone:8B2

Dilution WB 1:5000-1:10000, IHC 1:100-1:300, IF

1:100-1:300

Background

There are five tubulins in human cells: alpha, beta, gamma, delta, and epsilon. Tubulins are conserved across species. They form heterodimers, which multimerize to form a microtubule filament. An alpha and beta tubulin heterodimer is the basic structural unit of microtubules. The heterodimer does not come apart, once formed. The alpha and beta tubulins, which are each about 55 kDa MW, are homologous but not identical. Alpha, beta, and gamma tubulins have all been used as loading controls. Tubulin expression may vary according to resistance to antimicrobial and antimitotic drugs.

For Research Use Only

Thank you for your recent purchase

If you would like to learn more about antibodies, please visit www.elabscience.com.

Focus on your research Service for life science