


Tel:240-252-7368(USA) Fax: 240-252-7376(USA) techsupport@elabscience.com Website: www.elabscience.com

CDK4 Polyclonal Antibody


E-AB-52141 Catalog No. Reactivity H,M,R Storage Store at -20°C. Avoid freeze / thaw cycles. Host Rabbit **Applications** WB,IHC,ELISA **Isotype IgG**

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Images



Western blot analysis of Hela cell lysate using CDK4 Polyclonal Antibody at dilution of 1:1000



Immunohistochemistry of paraffinembedded Human esophagus cancer tissue using CDK4 Polyclonal Antibody at dilution of $1:60(\times 200)$

Immunogen Information

Fusion protein of human CDK4 **Immunogen**

BC003644 **Gene Accession** P11802 **Swissprot**

Synonyms Cdk 4,cdk4,CDK4 protein,CDK4,Cell division kinase

4,CMM 3,CMM3,p34 cdk4,PSK J3,PSK-J3

Product Information

Calculated MW 34 kDa

Observed MW Refer to figures

Buffer PBS with 0.05% NaN3 and 40% Glycerol,pH7.4

Purify Antigen affinity purification

Dilution WB 1:500-1:2000, IHC 1:50-1:300, ELISA

1:5000-1:10000

Background

The protein encoded by this gene is a member of the Ser/Thr protein kinase family. This protein is highly similar to the gene products of S. cerevisiae cdc28 and S. pombe cdc2. It is a catalytic subunit of the protein kinase complex that is important for cell cycle G1 phase progression. The activity of this kinase is restricted to the G1-S phase, which is controlled by the regulatory subunits D-type cyclins and CDK inhibitor p16(INK4a). This kinase was shown to be responsible for the phosphorylation of retinoblastoma gene product (Rb). Mutations in this gene as well as in its related proteins including D-type cyclins, p16(INK4a) and Rb were all found to be associated with tumorigenesis of a variety of cancers. Multiple polyadenylation sites of this gene have been reported.