

Tel:240-252-7368(USA) Fax: 240-252-7376(USA) techsupport@elabscience.com Website: www.elabscience.com

PPM1D Polyclonal Antibody


Catalog No.E-AB-52172ReactivityH,MStorageStore at -20°C. Avoid freeze / thaw cycles.HostRabbitApplicationsIHC,ELISAIsotypeIgG

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Images



Immunohistochemistry of paraffinembedded Human brain tissue using PPM1D Polyclonal Antibody at dilution of 1:60(×200)



Immunohistochemistry of paraffinembedded Human ovarian cancer tissue using PPM1D Polyclonal Antibody at dilution of 1:60(×200)

Immunogen Information

Immunogen Fusion protein of human PPM1D

Gene Accession BC033893 **Swissprot** O15297

Synonyms PP2C delta,PP2C-

delta,PP2CD,Ppm1d,PPM1D,Protein phosphatase magnesium-dependent 1 delta,Wild type p53 induced

phosphatase 1,WIP 1,WIP1

Product Information

Buffer PBS with 0.05% NaN3 and 40% Glycerol,pH7.4

Purify Antigen affinity purification

Dilution IHC 1:50-1:300, ELISA 1:2000-1:5000

Background

The protein encoded by this gene is a member of the PP2C family of Ser/Thr protein phosphatases. PP2C family members are known to be negative regulators of cell stress response pathways. The expression of this gene is induced in a p53-dependent manner in response to various environmental stresses. While being induced by tumor suppressor protein TP53/p53, this phosphatase negatively regulates the activity of p38 MAP kinase, MAPK/p38, through which it reduces the phosphorylation of p53, and in turn suppresses p53-mediated transcription and apoptosis. This phosphatase thus mediates a feedback regulation of p38-p53 signaling that contributes to growth inhibition and the suppression of stress induced apoptosis. This gene is located in a chromosomal region known to be amplified in breast cancer. The amplification of this gene has been detected in both breast cancer cell line and primary breast tumors, which suggests a role of this gene in cancer development.