


Tel:240-252-7368(USA) Fax: 240-252-7376(USA) techsupport@elabscience.com Website: www.elabscience.com

# **HIST1H1T Polyclonal Antibody**

E-AB-52289 Catalog No. Reactivity H,M Store at -20°C. Avoid freeze / thaw cycles. Rabbit **Storage** Host **Applications** WB,ELISA **Isotype IgG** 

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.

### **Images**



Western blot analysis of Mouse lung tissue using HIST1H1T Polyclonal Antibody at dilution of 1:400

## **Immunogen Information**

Full length fusion protein **Immunogen** 

**Gene Accession** BC130350 P22492 **Swissprot** 

**Synonyms** H1 histone family, member T (testis specific),H1FT

,H1T ,H1T,Hist1h1t,Histone cluster 1, H1t ,Histone

H1t, Testicular H1 histone

#### **Product Information**

Calculated MW 22 kDa

**Observed MW** Refer to figures

**Buffer** PBS with 0.05% NaN3 and 40% Glycerol,pH7.4

**Purify** Antigen affinity purification

**Dilution** WB 1:500-1:2000, ELISA 1:5000-1:10000

### **Background**

Histones are basic nuclear proteins responsible for nucleosome structure of the chromosomal fiber in eukaryotes. Two molecules of each of the four core histones (H2A, H2B, H3, and H4) form an octamer, around which approximately 146 bp of DNA is wrapped in repeating units, called nucleosomes. The linker histone, H1, interacts with linker DNA between nucleosomes and functions in the compaction of chromatin into higher order structures. This gene is intronless and encodes a replicationdependent histone that is a member of the histone H1 family.