


Tel:240-252-7368(USA) Fax: 240-252-7376(USA) techsupport@elabscience.com Website: www.elabscience.com

VASP Polyclonal Antibody

E-AB-52416 Catalog No. Reactivity H,M Storage Store at -20°C. Avoid freeze / thaw cycles. Rabbit Host **Applications** WB,IHC,ELISA **Isotype IgG**

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Images



Western blot analysis of A549 and HEPG2 cell lysates using VASP Polyclonal Antibody at dilution of 1:400



Immunohistochemistry of paraffinembedded Human tonsil tissue using VASP Polyclonal Antibody at dilution of $1:30(\times 200)$

Immunogen Information

Fusion protein of human VASP **Immunogen**

BC038224 **Gene Accession** P50552 **Swissprot**

Synonyms Vasodilator stimulated phosphoprotein, Vasodilator-

stimulated phosphoprotein, VASP, VASP

Product Information

Calculated MW 40 kDa

Observed MW Refer to figures

Buffer PBS with 0.05% NaN3 and 40% Glycerol,pH7.4

Purify Antigen affinity purification

Dilution WB 1:500-1:2000, IHC 1:25-1:100, ELISA

1:5000-1:10000

Background

Vasodilator-stimulated phosphoprotein (VASP) is a member of the Ena-VASP protein family. Ena-VASP family members contain an EHV1 Nterminal domain that binds proteins containing E/DFPPPXD/E motifs and targets Ena-VASP proteins to focal adhesions. In the mid-region of the protein, family members have a proline-rich domain that binds SH3 and WW domain-containing proteins. Their C-terminal EVH2 domain mediates tetramerization and binds both G and F actin. VASP is associated with filamentous actin formation and likely plays a widespread role in cell adhesion and motility. VASP may also be involved in the intracellular signaling pathways that regulate integrin-extracellular matrix interactions. VASP is regulated by the cyclic nucleotide-dependent kinases PKA and PKG.