


Tel:240-252-7368(USA) Fax: 240-252-7376(USA) techsupport@elabscience.com Website: www.elabscience.com

# PHF21A Polyclonal Antibody


Catalog No.E-AB-52472ReactivityH,MStorageStore at -20°C. Avoid freeze / thaw cycles.HostRabbitApplicationsIHC,ELISAIsotypeIgG

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.

# **Images**



Immunohistochemistry of paraffinembedded Human colorectal cancer tissue using PHF21A Polyclonal Antibody at dilution of 1:50(×200)



Immunohistochemistry of paraffinembedded Human ovarian cancer tissue using PHF21A Polyclonal Antibody at dilution of 1:50(×200)

# **Immunogen Information**

**Immunogen** Fusion protein of human PHF21A

**Gene Accession** BC015714 **Swissprot** Q96BD5

**Synonyms** BHC80a,BM-006,BRAF35-HDAC complex protein

BHC80,BRAF35/HDAC2 complex (80 kDa),KIAA1696,PF21A,PHD finger protein

21A,PHF21A

#### **Product Information**

**Buffer** PBS with 0.05% NaN3 and 40% Glycerol,pH7.4

**Purify** Antigen affinity purification

**Dilution** IHC 1:40-1:200, ELISA 1:5000-1:10000

# **Background**

Zinc-finger proteins contain DNA-binding domains and have a wide variety of functions, most of which encompass some form of transcriptional activation or repression. PHF21A (PHD finger protein 21A), also known as BRAF35-HDAC complex protein BHC80, is a 680 amino acid nuclear protein that contains one PHD-type zinc finger and one A.T hook DNA-binding domain, suggesting involvement in transcriptional regulation events. PHF21A is a component of the BHC complex, which is responsible for repressing transcription of neuron-specific genes in non-neuronal cells. The BHC complex acts as a chromatin modifier that deacetylates and demethylates specific sites on histones. PHF21A may act as a scaffold within the BHC complex. Predominantly expressed in brain, three isoforms of PHF21A exist as a result of alternative splicing events.