


Tel:240-252-7368(USA) Fax: 240-252-7376(USA) techsupport@elabscience.com Website: www.elabscience.com

PDIA4 Polyclonal Antibody

Catalog No.E-AB-52766ReactivityH,M,RStorageStore at -20°C. Avoid freeze / thaw cycles.HostRabbitApplicationsWB,IHC,ELISAIsotypeIgG

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Images



Western blot analysis of NIH/3T3 and RAW264.7 cell lysates using PDIA4 Polyclonal Antibody at dilution of 1:300



Immunohistochemistry of paraffinembedded Human cervical cancer tissue using PDIA4 Polyclonal Antibody at dilution of 1:50(×200)



Immunohistochemistry of paraffinembedded Human thyroid cancer tissue using PDIA4 Polyclonal Antibody at dilution of 1:50(×200)

Immunogen Information

Immunogen Fusion protein of human PDIA4

Gene Accession BC000425 **Swissprot** P13667

Synonyms EPR70,EPR72,ER protein 72,ERP 70,ERp72,PDIA

4,Pdia4,PDIA4,PDIR,Protein ERp72

Product Information

Calculated MW 73 kDa

Observed MW Refer to figures

Buffer PBS with 0.05% NaN3 and 40% Glycerol,pH7.4

Purify Antigen affinity purification

Dilution WB 1:500-1:2000, IHC 1:30-1:150, ELISA

1:5000-1:10000

Background

PDIA4(Protein disulfide-isomerase A4) is also named as ERP70,ERP72 and belongs to the protein disulfide isomerase family. It catalyzes the rearrangement of -S-S- bonds in proteins. ERp72 is a soluble protein localized in the ER lumen and contains the COOH-terminal retention signal,KEEL. There are 6 cysteine residues in the amino acid sequences of mouse and human ERp72. All of the cysteine residues occur in the internal thioredoxin motif,CGHC. The full length protein has a signal peptide with 20 amino acid.

For Research Use Only

Thank you for your recent purchase.

If you would like to learn more about antibodies, please visit www.elabscience.com.

Focus on your research Service for life science