


Tel:240-252-7368(USA) Fax: 240-252-7376(USA) techsupport@elabscience.com Website: www.elabscience.com

ECH1 Polyclonal Antibody


Catalog No.E-AB-52890ReactivityH,RStorageStore at -20°C. Avoid freeze / thaw cycles.HostRabbitApplicationsWB,IHC,ELISAIsotypeIgG

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Images



Western blot analysis of 293T and Hela cell Human fetal liver tissue lysates using ECH1 Polyclonal Antibody at dilution of 1:300



Immunohistochemistry of paraffinembedded Human thyroid cancer tissue using ECH1 Polyclonal Antibody at dilution of 1:50(×200)

Immunogen Information

Immunogen Fusion protein of human ECH1

Gene Accession BC011792 **Swissprot** Q13011

Synonyms ECH 1,Ech1,ECH1,HPXEL,mitochondrial,MS1034,P

eroxisomal enoyl CoA hydratase 1

Product Information

Calculated MW 36 kDa

Observed MW Refer to figures

Buffer PBS with 0.05% NaN3 and 40% Glycerol,pH7.4

Purify Antigen affinity purification

Dilution WB 1:500-1:2000, IHC 1:50-1:300, ELISA

1:5000-1:10000

Background

This gene encodes a member of the hydratase/isomerase superfamily. The gene product shows high sequence similarity to enoyl-coenzyme A (CoA) hydratases of several species, particularly within a conserved domain characteristic of these proteins. The encoded protein, which contains a C-terminal peroxisomal targeting sequence, localizes to the peroxisome. The rat ortholog, which localizes to the matrix of both the peroxisome and mitochondria, can isomerize 3-trans,5-cis-dienoyl-CoA to 2-trans,4-trans-dienoyl-CoA, indicating that it is a delta3,5-delta2,4-dienoyl-CoA isomerase. This enzyme functions in the auxiliary step of the fatty acid beta-oxidation pathway. Expression of the rat gene is induced by peroxisome proliferators.