


Tel:240-252-7368(USA) Fax: 240-252-7376(USA) techsupport@elabscience.com Website: www.elabscience.com

DUT Polyclonal Antibody


Catalog No.E-AB-52896ReactivityH,RStorageStore at -20°C. Avoid freeze / thaw cycles.HostRabbitApplicationsIHC,ELISAIsotypeIgG

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Images



Immunohistochemistry of paraffinembedded Human liver cancer tissue using DUT Polyclonal Antibody at dilution of 1:60(×200)



Immunohistochemistry of paraffinembedded Human thyroid cancer tissue using DUT Polyclonal Antibody at dilution of 1:60(×200)

Immunogen Information

Immunogen Fusion protein of human DUT

Gene Accession BC033645 **Swissprot** P33316

Synonyms nuclear isoform, mitochondrial, Deoxyuridine

triphosphatase, dut, DUT, dUTPase, FLJ20622

Product Information

Buffer PBS with 0.05% NaN3 and 40% Glycerol,pH7.4

Purify Antigen affinity purification

Dilution IHC 1:50-1:300, ELISA 1:5000-1:10000

Background

This gene encodes an essential enzyme of nucleotide metabolism. The encoded protein forms a ubiquitous, homotetrameric enzyme that hydrolyzes dUTP to dUMP and pyrophosphate. This reaction serves two cellular purposes: providing a precursor (dUMP) for the synthesis of thymine nucleotides needed for DNA replication, and limiting intracellular pools of dUTP. Elevated levels of dUTP lead to increased incorporation of uracil into DNA, which induces extensive excision repair mediated by uracil glycosylase. This repair process, resulting in the removal and reincorporation of dUTP, is self-defeating and leads to DNA fragmentation and cell death. Alternative splicing of this gene leads to different isoforms that localize to either the mitochondrion or nucleus. A related pseudogene is located on chromosome 19.