


Tel:240-252-7368(USA) Fax: 240-252-7376(USA) techsupport@elabscience.com Website: www.elabscience.com

# **PEPD Polyclonal Antibody**


Catalog No.E-AB-52976ReactivityH,M,RStorageStore at -20°C. Avoid freeze / thaw cycles.HostRabbitApplicationsWB,IHC,ELISAIsotypeIgG

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


# **Images**



Western blot analysis of HepG2 cell Mouse small intestines tissue lysates using PEPD Polyclonal Antibody at dilution of 1:1000



Immunohistochemistry of paraffinembedded Human liver cancer tissue using PEPD Polyclonal Antibody at dilution of 1:85(×200)



Immunohistochemistry of paraffinembedded Human tonsil tissue using PEPD Polyclonal Antibody at dilution of 1:85(×200)

## **Immunogen Information**

Immunogen Fusion protein of human PEPD

**Gene Accession** BC028295 **Swissprot** P12955

Synonyms MGC10905,MGC95081,Pep

4,Pep4,pepD,PEPD,Peptidase 4,Peptidase D,Prolidase

#### **Product Information**

Calculated MW 55 kDa

**Observed MW** Refer to figures

**Buffer** PBS with 0.05% NaN3 and 40% Glycerol,pH7.4

**Purify** Antigen affinity purification

**Dilution** WB 1:1000-1:5000, IHC 1:50-1:300, ELISA

1:5000-1:10000

### **Background**

This gene encodes a member of the peptidase family. The protein forms a homodimer that hydrolyzes dipeptides or tripeptides with C-terminal proline or hydroxyproline residues. The enzyme serves an important role in the recycling of proline, and may be rate limiting for the production of collagen. Mutations in this gene result in prolidase deficiency, which is characterized by the excretion of large amount of di- and tri-peptides containing proline. Multiple transcript variants encoding different isoforms have been found for this gene.

For Research Use Only

Thank you for your recent purchase

If you would like to learn more about antibodies, please visit www.elabscience.com.

Focus on your research Service for life science