


Tel:240-252-7368(USA) Fax: 240-252-7376(USA) techsupport@elabscience.com Website: www.elabscience.com

H2AFJ Polyclonal Antibody


E-AB-53031 Reactivity Catalog No. H,M,R Storage Store at -20°C. Avoid freeze / thaw cycles. Rabbit Host **Applications** WB,IHC,ELISA **Isotype IgG**

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Images



Western blot analysis of Mouse brain tissue 231 Jurkat cell lysates using H2AFJ Polyclonal Antibody at dilution of 1:800



Immunohistochemistry of paraffinembedded Human esophagus cancer tissue using H2AFJ Polyclonal Antibody at dilution of $1:160(\times 200)$

Immunogen Information

Fusion protein of human H2AFJ **Immunogen**

BC003602 **Gene Accession** O9BTM1 **Swissprot**

Synonyms H2A Histone Family Member J, Histone H2A. J, H2a/J

Product Information

Calculated MW 14 kDa

Observed MW Refer to figures

Buffer PBS with 0.05% NaN3 and 40% Glycerol,pH7.4

Purify Antigen affinity purification

Dilution WB 1:500-1:2000, IHC 1:150-1:300, ELISA

1:5000-1:10000

Background

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Nucleosomes consist of approximately 146 bp of DNA wrapped around a histone octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is located on chromosome 12 and encodes a replication-independent histone that is a variant H2A histone. The protein is divergent at the C-terminus compared to the consensus H2A histone family member. This gene also encodes an antimicrobial peptide with antibacterial and antifungal activity.