


Tel:240-252-7368(USA) Fax: 240-252-7376(USA) techsupport@elabscience.com Website: www.elabscience.com

# **SKP2 Polyclonal Antibody**


Catalog No.E-AB-65710ReactivityH,M,RStorageStore at -20°C. Avoid freeze / thaw cycles.HostRabbitApplicationsWB,IFIsotypeIgG

Note: Centrifuge before opening to ensure complete recovery of vial contents.

# **Images**



Western blot analysis of extracts of various cell lines using SKP2 Polyclonal Antibody at 1:500 dilution.



Immunofluorescence analysis of U2OS cells using SKP2 Polyclonal Antibody at dilution of 1:100. Blue: DAPI for nuclear staining.

# **Immunogen Information**

**Immunogen** A synthetic peptide of human SKP2

**GeneID** 6502 **Swissprot** Q13309

Synonyms SKP2,FBL1,FBXL1,FLB1,p45

#### **Product Information**

Calculated MW 23kDa/46kDa/47kDa

**Observed MW** 48KDa

**Buffer** PBS with 0.05% proclin300,50% glycerol,pH7.3.

**Purify** Affinity purification

**Dilution** WB 1:500-1:2000,IF 1:50-1:200

# **Background**

This gene encodes a member of the F-box protein family which is characterized by an approximately 40 amino acid motif, the F-box. The Fbox proteins constitute one of the four subunits of ubiquitin protein ligase complex called SCFs (SKP1-cullin-F-box), which function in phosphorylation-dependent ubiquitination. The F-box proteins are divided into 3 classes: Fbws containing WD-40 domains, Fbls containing leucinerich repeats, and Fbxs containing either different protein-protein interaction modules or no recognizable motifs. The protein encoded by this gene belongs to the Fbls class; in addition to an F-box, this protein contains 10 tandem leucine-rich repeats. This protein is an essential element of the cyclin A-CDK2 S-phase kinase. It specifically recognizes phosphorylated cyclin-dependent kinase inhibitor 1B (CDKN1B, also referred to as p27 or KIP1) predominantly in S phase and interacts with Sphase kinase-associated protein 1 (SKP1 or p19). In addition, this gene is established as a protooncogene causally involved in the pathogenesis of lymphomas. Alternative splicing of this gene generates three transcript variants encoding different isoforms.