


Tel:240-252-7368(USA) Fax: 240-252-7376(USA) techsupport@elabscience.com Website: www.elabscience.com

HLA-DPB1 Polyclonal Antibody

Catalog No.E-AB-66046ReactivityH,M,RStorageStore at -20°C. Avoid freeze / thaw cycles.HostRabbitApplicationsIHCIsotypeIgG

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Images



Immunohistochemistry of paraffinembedded Rat ovary using HLA-DPB1 Polyclonal Antibody at dilution of 1:100 (40x lens).



Immunohistochemistry of paraffinembedded Human colon carcinoma using HLA-DPB1 Polyclonal Antibody at dilution of 1:100 (40x lens).

Immunogen Information

Immunogen Recombinant fusion protein of human HLA-DPB1

(NP_002112.3).

GeneID 3115 **Swissprot** P04440

Synonyms DPB1,HLA-DP,HLA-DP1B,HLA-DPB,HLA-

DPB1,major histocompatibility complex

Product Information

Buffer PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Purify Affinity purification **Dilution** IHC 1:50-1:200

Background

HLA-DPB belongs to the HLA class II beta chain paralogues. This class II molecule is a heterodimer consisting of an alpha (DPA) and a beta chain (DPB), both anchored in the membrane. It plays a central role in the immune system by presenting peptides derived from extracellular proteins. Class II molecules are expressed in antigen presenting cells (APC: B lymphocytes, dendritic cells, macrophages). The beta chain is approximately 26-28 kDa and its gene contains 6 exons. Exon one encodes the leader peptide, exons 2 and 3 encode the two extracellular domains, exon 4 encodes the transmembrane domain and exon 5 encodes the cytoplasmic tail. Within the DP molecule both the alpha chain and the beta chain contain the polymorphisms specifying the peptide binding specificities, resulting in up to 4 different molecules.