


Tel:240-252-7368(USA) Fax: 240-252-7376(USA) techsupport@elabscience.com Website: www.elabscience.com

# CD68 Polyclonal Antibody


E-AB-66399 Reactivity Catalog No. H,M Storage Store at -20°C. Avoid freeze / thaw cycles. Rabbit Host **Applications** IHC,IF **Isotype IgG** 

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.

## **Images**



Immunohistochemistry of paraffinembedded Human liver using CD68 Polyclonal Antibody at dilution of 1:200 (40x lens).



Immunofluorescence analysis of RAW264.7 cells using CD68 Polyclonal Antibody at dilution of 1:100. Blue: DAPI for nuclear staining.

### **Immunogen Information**

**Immunogen** Recombinant fusion protein of human CD68

(NP\_001242.2).

GeneID 968 **Swissprot** P34810

**Synonyms** CD68,GP110,LAMP4,SCARD1

#### **Product Information**

**Buffer** PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Purify Affinity purification

**Dilution** IHC 1:50-1:200 IF 1:50-1:200

## **Background**

This gene encodes a 110-kD transmembrane glycoprotein that is highly expressed by human monocytes and tissue macrophages. It is a member of the lysosomal/endosomal-associated membrane glycoprotein (LAMP) family. The protein primarily localizes to lysosomes and endosomes with a smaller fraction circulating to the cell surface. It is a type I integral membrane protein with a heavily glycosylated extracellular domain and binds to tissue- and organ-specific lectins or selectins. The protein is also a member of the scavenger receptor family. Scavenger receptors typically function to clear cellular debris, promote phagocytosis, and mediate the recruitment and activation of macrophages. Alternative splicing results in multiple transcripts encoding different isoforms.