


Tel:240-252-7368(USA) Fax: 240-252-7376(USA) techsupport@elabscience.com Website: www.elabscience.com

c-Myc Polyclonal Antibody

E-AB-67107 Catalog No. Reactivity H,M Storage Store at -20°C. Avoid freeze / thaw cycles. Rabbit Host **Applications Isotype IgG**

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Images



Western blot analysis of extracts of various cell lines using c-Myc Polyclonal Antibody at dilution of 1:1000.

Immunogen Information

A synthetic peptide of human c-Myc **Immunogen**

4609 GeneID **Swissprot** P01106

Synonyms MRTL, MYCC, bHLHe39, c-Myc, MYC

Product Information

Calculated MW 48kDa/50kDa

Observed MW 65kDa

Buffer PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Purify Affinity purification Dilution WB 1:500-1:2000

Background

The protein encoded by this gene is a multifunctional, nuclear phosphoprotein that plays a role in cell cycle progression, apoptosis and cellular transformation. It functions as a transcription factor that regulates transcription of specific target genes. Mutations, overexpression, rearrangement and translocation of this gene have been associated with a variety of hematopoietic tumors, leukemias and lymphomas, including Burkitt lymphoma. There is evidence to show that alternative translation initiations from an upstream, in-frame non-AUG (CUG) and a downstream AUG start site result in the production of two isoforms with distinct N-termini. The synthesis of non-AUG initiated protein is suppressed in Burkitt's lymphomas, suggesting its importance in the normal function of this gene.