


Tel:240-252-7368(USA) Fax: 240-252-7376(USA) techsupport@elabscience.com Website: www.elabscience.com

# **CCNT1 Polyclonal Antibody**

Catalog No. E-AB-67400 Reactivity Η Store at -20°C. Avoid freeze / thaw cycles. Rabbit **Storage** Host **Applications Isotype IgG** 

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.

## **Images**



Western blot analysis of extracts of K-562 cells using CCNT1 Polyclonal Antibody.

## **Immunogen Information**

**Immunogen** A synthetic peptide of human CCNT1

(NP\_001231.2).

GeneID 904 **Swissprot** O60563

**Synonyms** CCNT1,CCNT,CYCT1,HIVE1,cyclin-T1

#### **Product Information**

**Calculated MW** 21kDa/80kDa

**Observed MW** 87kDa

**Buffer** PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Purify** Affinity purification Dilution WB 1:500-1:2000

## **Background**

This gene encodes a member of the highly conserved cyclin C subfamily. The encoded protein tightly associates with cyclin-dependent kinase 9, and is a major subunit of positive transcription elongation factor b (p-TEFb). In humans, there are multiple forms of positive transcription elongation factor b, which may include one of several different cyclins along with cyclin-dependent kinase 9. The complex containing the encoded cyclin and cyclin-dependent kinase 9 acts as a cofactor of human immunodeficiency virus type 1 (HIV-1) Tat protein, and is both necessary and sufficient for full activation of viral transcription. This cyclin and its kinase partner are also involved in triggering transcript elongation through phosphorylation of the carboxy-terminal domain of the largest RNA polymerase II subunit. Overexpression of this gene is implicated in tumor growth. Alternative splicing results in multiple transcript variants.