


Tel:240-252-7368(USA) Fax: 240-252-7376(USA) techsupport@elabscience.com Website: www.elabscience.com

DCTN2 Polyclonal Antibody


Catalog No. E-AB-67423 Reactivity H,R Store at -20°C. Avoid freeze / thaw cycles. Rabbit **Storage** Host **Applications** WB.IF **Isotype IgG**

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Images



Western blot analysis of extracts of MCF-7 cells using DCTN2 Polyclonal Antibody at 1:1000 dilution.



Immunofluorescence analysis of U2OS cells using DCTN2 Polyclonal Antibody Blue: DAPI for nuclear staining.

Immunogen Information

Immunogen Recombinant fusion protein of human DCTN2

GeneID 10540 **Swissprot** O13561

Synonyms DCTN2.DCTN50.DYNAMITIN.HEL-S-77.RBP50

Product Information

Calculated MW 44kDa **Observed MW** 44kDa

Buffer PBS with 0.02% sodium azide,50% glycerol,pH7.3.

Purify Affinity purification

Dilution WB 1:500-1:2000,IF 1:20-1:50

Background

This gene encodes a 50-kD subunit of dynactin, a macromolecular complex consisting of 10-11 subunits ranging in size from 22 to 150 kD. Dynactin binds to both microtubules and cytoplasmic dynein. It is involved in a diverse array of cellular functions, including ER-to-Golgi transport, the centripetal movement of lysosomes and endosomes, spindle formation, chromosome movement, nuclear positioning, and axonogenesis. This subunit is present in 4-5 copies per dynactin molecule. It contains three short alpha-helical coiled-coil domains that may mediate association with self or other dynactin subunits. It may interact directly with the largest subunit (p150) of dynactin and may affix p150 in place. Multiple alternatively spliced transcript variants encoding distinct isoforms have been found for this gene.