


Tel:240-252-7368(USA) Fax: 240-252-7376(USA) techsupport@elabscience.com Website: www.elabscience.com

# (KO Validated) MTA2 Polyclonal Antibody

Catalog No.E-AB-67426ReactivityH,M,RStorageStore at -20°C. Avoid freeze / thaw cycles.HostRabbitApplicationsWBIsotypeIgG

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.

# **Images**



Western blot analysis of extracts of various cell lines using MTA2 Polyclonal Antibody.

# **Immunogen Information**

**Immunogen** Recombinant fusion protein of human MTA2

(NP\_004730.2).

**GeneID** 9219 **Swissprot** 094776

Synonyms MTA2,MTA1L1,PID

#### **Product Information**

Calculated MW 55kDa/75kDa

Observed MW 72kDa

**Buffer** PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Purify Affinity purification
Dilution WB 1:500-1:2000

# **Background**

This gene encodes a protein that has been identified as a component of NuRD, a nucleosome remodeling deacetylase complex identified in the nucleus of human cells. It shows a very broad expression pattern and is strongly expressed in many tissues. It may represent one member of a small gene family that encode different but related proteins involved either directly or indirectly in transcriptional regulation. Their indirect effects on transcriptional regulation may include chromatin remodeling. It is closely related to another member of this family, a protein that has been correlated with the metastatic potential of certain carcinomas. These two proteins are so closely related that they share the same types of domains. These domains include two DNA binding domains, a dimerization domain, and a domain commonly found in proteins that methylate DNA. One of the proteins known to be a target protein for this gene product is p53. Deacetylation of p53 is correlated with a loss of growth inhibition in transformed cells supporting a connection between these gene family members and metastasis.